



Drivers of antibiotic prescribing in children and adolescents with febrile lower respiratory tract infections

Gotta, Verena ; Baumann, Philipp ; Ritz, Nicole ; Fuchs, Aline ; Baer, Gurli ; Bonhoeffer, Jessica M ; Heininger, Ulrich ; Szinnai, Gabor ; Bonhoeffer, Jan ; ProPAED study group

Abstract: **BACKGROUND:** Knowledge of key drivers for antibiotic prescribing in pediatric lower respiratory tract infection (LRTI) could support rational antibiotic use. Thus, we aimed to determine the impact of clinical and laboratory factors on antibiotic prescribing in children and adolescents with febrile LRTI. **METHODS:** Pediatric patients from the standard care control group of a randomized controlled trial (ProPAED) investigating procalcitonin guided antibiotic treatment in febrile LRTI were included in a multivariate logistic regression analysis to evaluate the impact of laboratory and clinical factors on antibiotic prescribing. **RESULTS:** The standard care control group of the ProPAED study comprised 165 LRTI patients (median age: 2.7 years, range: 0.1-16), out of which 88 (55%) received antibiotic treatment. Factors significantly associated with antibiotic prescribing in patients with complete clinical and laboratory documentation ($n = 158$) were C-reactive protein (OR 5.8 for a 10-fold increase, 95%CI 2.2-14.9), white blood count beyond age-dependent reference range (OR 3.9, 95%CI 1.4-11.4), body temperature (OR 1.7 for an increase by 1°C, 95%CI 1.02-2.68), and pleuritic pain (OR 2.8, 95%CI 1.1-7.6). Dyspnea (OR 0.3, 95%CI 0.1-0.7) and wheezing (OR 0.3, 95%CI 0.13-0.95) were inversely associated with antibiotic prescribing. **CONCLUSION:** Laboratory markers were strong drivers of antibiotic prescribing in children with febrile lower respiratory tract infections, in spite of their known poor prediction of antibiotic need. Building on current guidelines for antibiotic treatment in children with febrile LRTI, a reliable decision algorithm for safe antibiotic withholding considering the laboratory and clinical factors evaluated in this study has the potential to further reduce antibiotic prescribing.

DOI: <https://doi.org/10.1371/journal.pone.0185197>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-146926>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Gotta, Verena; Baumann, Philipp; Ritz, Nicole; Fuchs, Aline; Baer, Gurli; Bonhoeffer, Jessica M; Heininger, Ulrich; Szinnai, Gabor; Bonhoeffer, Jan; ProPAED study group (2017). Drivers of antibi-

otic prescribing in children and adolescents with febrile lower respiratory tract infections. PLoS ONE, 12(9):e0185197.
DOI: <https://doi.org/10.1371/journal.pone.0185197>

RESEARCH ARTICLE

Drivers of antibiotic prescribing in children and adolescents with febrile lower respiratory tract infections

Verena Gotta¹✉, Philipp Baumann^{2,3,4,*}, Nicole Ritz^{1,2}, Aline Fuchs¹, Gurli Baer², Jessica M. Bonhoeffer^{3,4}, Ulrich Heininger², Gabor Szinnai⁴, Jan Bonhoeffer², for the ProPAED study group[¶]

1 Department of Pediatric Pharmacology and Pharmacometrics, University of Basel Children's Hospital, Basel, Switzerland, **2** Department of Pediatric Infectious Diseases and Vaccinology, University of Basel Children's Hospital, Basel, Switzerland, **3** University of Basel Children's Hospital, Basel, Switzerland, **4** Department of Pediatric Endocrinology and Diabetology, University of Basel Children's Hospital, Basel, Switzerland

✉ These authors contributed equally to this work.

✉ Current address: Department of Intensive Care Medicine and Neonatology, University Children's Hospital Zurich, Zurich, Switzerland

✉ Current address: Child Development Center, University Children's Hospital Zurich, Zurich, Switzerland

¶ Membership of the ProPAED study group is listed in the Acknowledgments.

* philipp.baumann@kispi.uzh.ch



OPEN ACCESS

Citation: Gotta V, Baumann P, Ritz N, Fuchs A, Baer G, Bonhoeffer JM, et al. (2017) Drivers of antibiotic prescribing in children and adolescents with febrile lower respiratory tract infections. PLoS ONE 12(9): e0185197. <https://doi.org/10.1371/journal.pone.0185197>

Editor: Jacobus P. van Wouwe, TNO, NETHERLANDS

Received: April 11, 2017

Accepted: September 5, 2017

Published: September 28, 2017

Copyright: © 2017 Gotta et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: The authors received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Abstract

Background

Knowledge of key drivers for antibiotic prescribing in pediatric lower respiratory tract infection (LRTI) could support rational antibiotic use. Thus, we aimed to determine the impact of clinical and laboratory factors on antibiotic prescribing in children and adolescents with febrile LRTI.

Methods

Pediatric patients from the standard care control group of a randomized controlled trial (ProPAED) investigating procalcitonin guided antibiotic treatment in febrile LRTI were included in a multivariate logistic regression analysis to evaluate the impact of laboratory and clinical factors on antibiotic prescribing.

Results

The standard care control group of the ProPAED study comprised 165 LRTI patients (median age: 2.7 years, range: 0.1–16), out of which 88 (55%) received antibiotic treatment. Factors significantly associated with antibiotic prescribing in patients with complete clinical and laboratory documentation (n = 158) were C-reactive protein (OR 5.8 for a 10-fold increase, 95%CI 2.2–14.9), white blood count beyond age-dependent reference range (OR 3.9, 95%CI 1.4–11.4), body temperature (OR 1.7 for an increase by 1°C, 95%CI 1.02–2.68), and pleuritic pain (OR 2.8, 95%CI 1.1–7.6). Dyspnea (OR 0.3, 95%CI 0.1–0.7) and wheezing (OR 0.3, 95%CI 0.13–0.95) were inversely associated with antibiotic prescribing.

Conclusion

Laboratory markers were strong drivers of antibiotic prescribing in children with febrile lower respiratory tract infections, in spite of their known poor prediction of antibiotic need. Building on current guidelines for antibiotic treatment in children with febrile LRTI, a reliable decision algorithm for safe antibiotic withholding considering the laboratory and clinical factors evaluated in this study has the potential to further reduce antibiotic prescribing.

Introduction

Lower respiratory tract infections (LRTI) bear high mortality and morbidity in children and adolescents with community acquired pneumonia (CAP) being the leading cause of death worldwide with 1.3 million fatal pediatric cases each year [1]. Further, LRTI are associated with non-fatal complications like pleural effusions, empyema (up to 12% of pneumonia cases) or necrotizing pneumonia (0.8%) which are feared by both hospital and general physicians [2, 3]. However, identifying patients benefitting from antibiotic treatment among those not in need remains a major diagnostic challenge. Thus, physicians tend to err on the safe side and overuse antibiotics according to varied approaches as reflected in the different antibiotic prescription rates across countries and institutions [4–8]. A better understanding of the decision to prescribe or withhold antibiotics is crucial for future restrictive treatment strategies. Factors leading to antibiotic prescription are known to be complex [5] and are scarcely elucidated in the pediatric population. Given that adult studies have identified e.g. a strong impact of C-reactive protein (CRP) (odds ratio 98.1 for exceeding 50 mg/L) on physicians' decision to prescribe antibiotic medication for LRTI [9] there is a need for clarification of what sways the decision to prescribe antibiotics for LRTI in children and adolescents. Thus, we aimed at characterizing the laboratory and clinical drivers of antibiotic prescribing for pediatric LRTI with a subanalysis of the ProPAED trial [10]. This study was among the first randomized controlled trials in children evaluating the feasibility and effect of Procalcitonin (PCT) guided treatment for respiratory tract infections [11, 12] after first observational studies suggested that PCT might be considered in pediatrics. The ProPAED investigators tested established adult PCT cut-off ranges in a cohort of pediatric emergency departments of tertiary care centers [13–19].

Methods

Study population

This is a hypothesis generating post-hoc analysis from a randomized controlled trial (ProPAED) evaluating PCT as a biomarker to reduce antibiotic treatment of LRTI in children and adolescents [10]. Briefly, children and adolescents 1 month to 18 years of age presenting with LRTI to the pediatric emergency departments of two pediatric tertiary care centers in Switzerland between 01/2009 and 02/2010 were included. Acute LRTI was defined by the presence of fever ($\geq 38^{\circ}\text{C}$ measured in the hospital at study enrolment or at home), and at least one of the following symptoms: tachypnea, dyspnea, wheezing, late inspiratory crackles, bronchial breathing, and/or pleural rub. CAP was defined as febrile LRTI with a new or increasing alveolar infiltrate on chest radiograph. In the original ProPAED trial patients were randomized to the PCT guided intervention group or to the standard care control group with treatment based on international guidelines [4]. For the analysis of factors associated with antibiotic prescribing only patients from the standard care control group were included. Follow-up was available

for 98% of all ProPAED patients and no increase of adverse events was associated with antibiotic withholding in either of the treatment groups [10].

Variables

The following variables were included for analysis. As clinical parameters presence or absence of pleuritic pain, pleural rub, wheezing, tachypnea, dyspnea, late inspiratory crackles, reduced breathing sound, bronchial breathing, body temperature, and heart and respiratory rate were chosen. For demographic variables and patient history age, pneumococcal and *Haemophilus influenzae* type b vaccination status (as classified according to S1 and S2 Tables), and days of fever preceding presentation at hospital were included. As laboratory and microbiology parameters CRP, and white blood cell count (WBC), respiratory syncytial virus (RSV) type A/B or influenza A/B on nasopharyngeal aspirate (NPA) were included. All laboratory and microbiological tests were performed at study enrolment. Blood culture results were not included in this analysis since they were not systematically obtained and therefore available only in 72 (43%) of patients with 4 positive results.

Definitions

WBC was classified as normal or beyond age-dependent reference range according to reference values provided by the hospital-specific laboratory (S3 and S4 Tables). Heart rate was classified as normal or elevated according to age-dependent reference ranges (S5 Table) [20], respiratory rate was classified as elevated if exceeding WHO age-dependent reference ranges for children up to 5 years of age, and if exceeding percentile 90 as reported by Fleming et al. for children >5 years of age (S6 Table) [21].

Statistical analysis

Analysis of factors associated with antibiotic prescription. The association of clinical, demographic and laboratory variables with antibiotic prescribing was assessed using exploratory graphical evaluation and univariate logistic regression analysis (S1 Fig). For variables showing a significant association, sensitivity, specificity, accuracy and area under the receiver operating characteristics (ROC) curve were calculated as performance measures to predict or classify antibiotic prescribing. For continuous variables these performance measures were calculated for the best threshold (Youden index) which was determined by ROC analysis. Subsequently, variables showing a significant association in univariate analysis were included in a multivariate logistic regression model, followed by stepwise backward deletion of non-significant variables. The predictive performance of the final multivariate model was expressed regarding accuracy, sensitivity and specificity (using a probability threshold of ≥ 0.5 for classification of antibiotic prescribing). Continuous variables were partly transformed (e.g. log-transformation, categorization) and centered to normal values (instead of zero) for regression analysis to linearize relationships, according to clinical relevance and graphical evaluation of the appropriateness of assumed relationships (S2–S5 Figs). Missing data was ignored (complete-case analysis), other than vaccination status and NPA test results, where missingness was interpreted as incomplete vaccination status or negative NPA test result, respectively. The global goodness of fit of the final multivariate model was assessed by unweighted sum of squares statistic (Cessie, van Houwelingen, Copas, Hosmer unweighted sum of squares test) and residual diagnostics (Pearson residuals were plotted and any points $> |3$ standard deviations| were investigated); overdispersion was assessed and deemed not to be present if the scale parameter (estimate of the residual mean square) was within the interval [0.85, 1.15]. The

relative goodness of fit compared to simpler models following single term deletion was assessed using the likelihood-ratio test.

Sensitivity analyses. The following three sensitivity analyses were performed: (1) The interaction between pleuritic pain and age (stratified by 2 years of age), was assessed, since pain is difficult to be assessed children < 2 years. (2) Elastic net regression was used to verify robustness of selected covariates. (3) Remaining missing variables (<3%) were multiply imputed by fully conditional specification using the R package mice and re-tested in the logistic regression analysis [22]. All statistical analyses were performed in R (version 3.2.1; R Development Core Team, Vienna, Austria, <http://www.r-project.org>). Example codes of the main analyses can be found in the supplemental data (S1 Text).

Ethics

Both the ethics committee of the University of Basel and Kanton Aargau approved the trial protocol. Written informed consent was obtained from all patients or their care takers. The trial was registered with the International Standard Randomized Controlled Trial Number Register (ISRCTN 17057980).

Results

Factors associated with antibiotic prescription

Out of 169 patients in the standard care group, a total of 165 patients with complete data sets for the current analyses were included. Of these, 93 (56%) received antibiotic treatment (Table 1). The majority of patients treated with an antibiotic ($n = 78$, i.e. 84%) received it on the first day of study inclusion.

Eleven of 19 assessed clinical and laboratory variables showed a significant association with antibiotic prescribing (Table 2). Of these, eight variables were associated with increased antibiotic prescription and three with reduced antibiotic prescription. Corresponding odds ratio estimates from logistic regression analyses are summarized in Table 2 for all tested clinical and laboratory variables and illustrated in Fig 1 together with the respective multivariate estimates. See complete sensitivity for univariate regression analysis in S7 Table.

Fig 2 presents sensitivities and specificities for all eleven variables. The highest sensitivity was calculated for the absence of the symptom wheezing (90%). However, specificity was low (51%). Best specificity (92%) was calculated for a normal white blood cell count. However, here sensitivity was low (50%).

For multivariate logistic regression analysis: 158 patients with complete clinical and laboratory findings of all variables were included (Table 3). Of those, 87 (55%) received an antibiotic. Six of the initially eleven significant variables were retained in this final model, after step-wise backwards-deletion of non-significant associations. Model diagnostics of goodness of fit and dispersion did not indicate model misspecification, or preference of simpler models. Multi- and univariate odds ratio estimates are contrasted in Fig 1.

The multivariate model explained the antibiotic prescribing in 126 patients (accuracy: 80%, 95%CI: 73–86%), with a correct prediction of antibiotic prescription of 83% (sensitivity, 95% CI: 76–86%) and a correct prediction of withholding antibiotic therapy of 76% (specificity; 95%CI: 69–82%). Characteristics of falsely predicted patients are summarized in S8 Table.

Sensitivity analyses

Subgroup analysis indicated that the association of pleuritic pain with antibiotic prescribing was reduced in younger children (in < 2 years OR = 1.1, $p = 0.896$, $n = 65$) compared to the

Table 1. Patients' characteristics.

	Antibiotic treatment	No antibiotic treatment
Total number of patients [n (%)]	93	72
received antibiotic at day 1	78 (84)	NA
received antibiotic > day 1	15 (16)	NA
antibiotic pretreatment	16 (17)	1 (1)
Clinical diagnosis (at inclusion) [n (%)]		
Bronchitis/Bronchiolitis	10 (11)	50 (69)
Bronchitis/Bronchiolitis + Pneumonia	24 (26)	17 (24)
Pneumonia	59 (63)	5 (7)
Radiologic diagnosis [n (%)]		
Bronchitis/Bronchiolitis	23 (25)	41 (57)
Bronchopneumonia	49 (53)	11 (15)
Lobar pneumonia	17 (18)	1 (1)
Radiography not performed	4 (4)	19 (26)
Age in years [median (range)]	3.9 (0.1–15.3)	1.9 (0.1–15.5)
< 5 years [n (%)]	57 (61.3)	60 (83.3)
< 2 years [n (%)]	26 (28.0)	39 (54.2)
Temperature [°C (range)]	38.6 (36.3–40.7), NA = 1	38.2 (35.8–39.9)
Preceding days of fever [days (range)]	3 (1–14), NA = 2	2 (1–12)
Heart rate [1/min (range)]	138 (84–192), NA = 3	141 (88–193), NA = 2
Respiratory rate [1/min (range)]	38 (16–80), NA = 2	40 (18–80), NA = 3
C-reactive protein [mg/L (range)]	37 (3–653), NA = 3	9 (3–144), NA = 1
White blood cell count [G/L (range)]	13.4 (3.3–54.8), NA = 3	9.5 (4.4–24.4)
beyond reference range [n (%)]	46 (50) 6 low, 40 elevated	9 (12) 4 low, 5 elevated
normal [n (%)]	44 (40)	63 (88)
Symptoms at inclusion [n (%)]		
Wheezing	9 (10)	37 (51)
Dyspnea	47 (51)	57 (79)
Reduced breathing sound	37 (40)	10 (14)
Pleuritic pain	36 (39)	15 (21)
Crackles	20 (22)	24 (33)
Bronchial breathing	16 (17)	7 (10)
Late inspiratory crackles	42 (45)	25 (35)
Pleural rub	2 (2)	0
Vaccination status [n (%)]		
Pneumococcal vaccination		
Incomplete or unknown	68 (73, NA = 10)	36 (50, NA = 6)
<i>Haemophilus influenzae</i> type b vaccination		
Incomplete or unknown	24 (26, NA = 9)	18 (25, NA = 3)
Nasopharyngeal aspirate [n (%)]		
Negative or not performed	40 (43, NA = 5)	29 (40, NA = 5)
Human metapneumovirus	13 (14)	5 (7)
Respiratory syncytial virus A/B	7/6 (14)	18/6 (33)
Influenza A/B	5/1 (6)	3/1 (6)
H1N1	5	2
P1V1	1	2
<i>Mycoplasma pneumoniae</i>	7	1
<i>Chlamydia pneumoniae</i>	1	-

(Continued)

Table 1. (Continued)

	Antibiotic treatment	No antibiotic treatment
Adenovirus	1	1
Combinations	6	4
Hospitalization [n (%)]	61 (66)	34 (47)
Education (mother) [n (%)]		
School	15 (17)	21 (30)
Apprenticeship	48 (54)	30 (43)
Technical college	14 (16)	8 (11)
University	12 (13)	11 (16)

NA: not applicable

<https://doi.org/10.1371/journal.pone.0185197.t001>

Table 2. Variables associated with antibiotic prescribing in univariate logistic regression.

Variable (definition of baseline)	Baseline antibiotic treatment % (95%CI)	Univariate OR (95% CI)	Definition of variable effect	Univariate p-value
C-reactive protein (10 mg/L)	40 (31–50)	9.3 (4.3–19.8)	for a 10 fold increase	9.2×10^{-9}
Wheezing (absence)	71 (62–78)	0.1 (0.0–0.2)	presence of symptom	6.1×10^{-8}
WBC (normal for age)	41 (32–51)	7.3 (3.3–16.5)	> or < age-dependent reference range	1.5×10^{-6}
Age (0 years)	32 (20–46)	1.4 (1.2–1.7)	for each additional year of age until 5 years	0.0002
Temperature (37°C)	34 (22–47)	2.0 (1.4–2.9)	for each additional degree (in °C)	0.0002
Dyspnea (absence)	75 (63–85)	0.3 (0.1–0.5)	presence of symptom	0.0002
Reduced breathing sound (absence)	47 (39–56)	4.1 (1.9–9.0)	presence of symptom	0.0004
Pneumococcal vaccination (complete for age)	41 (29–54)	2.7 (1.4–5.2)	Incomplete or unknown	0.0026
Preceding days of fever (1 day)	44 (32–54)	1.3 (1.1–1.5)	for each additional preceding day of fever	0.0029
Respiratory syncytial virus + (negative or unknown NPA test)	62 (53–70)	0.4 (0.2–0.8)	positive NPA test	0.011
Pleuritic pain (absence)	50 (41–59)	2.4 (1.2–4.9)	presence of symptom	0.015
Crackles (absence)	60 (51–69)	0.5 (0.3–1.1)	presence of symptom	0.090
Bronchial breathing (absence)	54 (46–62)	1.9 (0.7–5.0)	presence of symptom	0.17
Late inspiratory crackles (absence of symptom)	52 (42–62)	1.5 (0.8–2.9)	presence of symptom	0.18
Heart rate (normal for age)	51 (38–64)	1.4 (0.7–2.7)	High for age	0.34
Respiratory rate (normal for age)	53 (41–64)	1.3 (0.7–2.5)	High for age	0.39
Hib vaccination (complete for age)	56 (47–65)	1.0 (0.5–2.1)	Incomplete or unknown	0.91
Influenzae (negative or unknown NPA test)	56 (49–64)	1.0 (0.2–3.7)	positive NPA test	0.96
Pleural rub (absence)	56 (48–63)	>10.0 (0-infinity)	presence of symptom	0.99

Number of patients included in univariate analysis: n = 165, besides for C-reactive protein: n = 161, temperature: n = 164, heart rate: n = 160, respiratory rate: n = 160, and WBC: n = 16. NPA: nasopharyngeal aspirate, OR: odds ratio, WBC: white blood cell count. p-values were used to order variables in decreasing order of statistical significance.

<https://doi.org/10.1371/journal.pone.0185197.t002>




Fig 1. Associations of laboratory and clinical factors with antibiotic prescribing. Odds ratio estimates with 95% confidence intervals (95%CI) from univariate and multivariate logistic regression analysis. Open circles and dashed lines: univariate OR and 95%CI; variables are presented in decreasing order of strength of association with antibiotic prescription (p-value, details: see Table 2), grey circles indicate a non-significant association (p-value ≥ 0.05), the OR estimate for pleural rub (OR $>> 10$ with 95%CI ranging from 0-infinity, Table 1) is not illustrated. Black dots and solid lines: multivariate OR with 95%CI of variables remaining after backward deletion of non-significant variables. Vertical line: OR = 1 indicating no association with antibiotic prescription, ORs > 1 indicate an association with increased antibiotic prescription, ORs < 1 indicate an association with reduced antibiotic use. CRP: C-reactive protein. WBC: White blood cell count. Vacc.: Vaccine. RSV: Respiratory syncytial virus. HR: Heart rate. Hib: *Haemophilus influenzae* type b.

<https://doi.org/10.1371/journal.pone.0185197.g001>

OR in the full analysis (OR = 2.4, $p = 0.015$, $n = 165$), and older children only (OR = 2.7, $p = 0.037$ in patients ≥ 2 years, $n = 100$); including a formal interaction term for this age-dependent association reached however not statistical significance of pleuritic pain in neither group. Elastic net regression suggested lowest misclassification error (optimal regularization parameter lambda from 100-fold cross-validation) when 9 covariates are included in a multivariate model (CRP, wheezing, WBC, temperature, dyspnea, preceding days of fever, RSV+, pleuritic pain, crackles). A simpler/more regularized model with only 4 covariates (CRP, Wheezing, WBC, dyspnea) could be favored, when accepting a larger error, within one standard error of the minimum. Using a higher penalty ($\alpha = 1$, i.e. lasso) for covariable selection, the same 4 covariates would be selected (CRP, wheezing, WBC, dyspnea) to obtain lowest misclassification error.

Results of univariate regression analysis did not significantly change when estimating coefficients from multiply imputed datasets (the same 11 variables showed significant associations with antibiotic prescribing with a p-value < 0.05). The multivariate regression results changed slightly, with pleuritic pain removed in backward deletion due to a p-value of 0.056 (compared to $p = 0.038$ in the initial analysis); all other 5 variables (CRP, wheezing, WBC, dyspnea, temperature) were retained with a p-value < 0.05 .

Discussion

This hypothesis generating study investigated associations of laboratory and clinical findings with antibiotic prescription in children and adolescents with LRTI presenting to the pediatric




Fig 2. Sensitivities and specificities for associations with antibiotic prescribing. Illustration of sensitivities and specificities (dots) with 95% confidence intervals (crosses) of each single variable for antibiotic prescribing. Left panel: dichotomous variables. Right panel: For continuous variables the range of sensitivities and specificities for all possible thresholds is illustrated (gray receiver operating characteristic curves), as well as the sensitivity and specificity associated with the best threshold estimates. Best threshold estimates (95%CI) were: CRP: 26 (12–55) mg/L, age: 2.1 (1.1–4.8) years, temperature: 38.6 (37.3–38.9) °C, preceding fever: 2.5 (1.5–5.5) days. CRP: C-reactive protein. RSV: Respiratory syncytial virus. WBC: White blood cell count.

<https://doi.org/10.1371/journal.pone.0185197.g002>

emergency department. The main finding of this study was the strong association of elevated routine laboratory parameters (CRP and WBC) with increased antibiotic prescribing. Four clinical factors (dyspnea, body temperature, pleuritic pain and wheezing) were also associated with antibiotic prescribing and explained, together with the laboratory parameters, 80% of the antibiotic prescription behavior.

Laboratory parameters associated with increased antibiotic prescription

Although the diagnostic utility of the laboratory values is known to be limited, in this study they have been found to play a major role for antibiotic prescribing. The strongest association with

Table 3. Variables associated with antibiotic prescription in multivariate logistic regression.

Variable (definition of baseline ^a)	Effect ^b (multivariate OR)	Definition of effect	p-value
C-reactive protein (10 mg/L)	5.8 (2.2–14.9)	for a 10 fold increase	0.0003
Dyspnea (absence of symptom)	0.3 (0.1–0.7)	presence of symptom	0.008
WBC (normal)	3.9 (1.4–11.4)	> or < age-dependent reference range	0.011
Temperature (37°C)	1.7 (1.0–2.7)	for each additional degree (in °C)	0.040
Pleuritic pain (absence of symptom)	2.8 (1.1–7.6)	presence of symptom	0.038
Wheezing (absence)	0.3 (0.1–0.9)	presence of symptom	0.039

^aBaseline probability estimate: 40% antibiotic prescription (20–64%).

^bVariables removed after stepwise backward deletion of non-significant associations (OR and p-values from full multivariate model): age (OR = 0.97, p = 0.84), reduced breathing sound (OR = 1.08, p = 0.89), incomplete pneumococcal vaccination status (OR = 0.94, p = 0.91), positive RSV test (OR = 0.51, p = 0.20), days of fever (OR = 1.20, p = 0.08). OR: Odds ratio. WBC: White blood cell count.

<https://doi.org/10.1371/journal.pone.0185197.t003>

prescription of antimicrobial agents was found in the multivariate analysis for higher than normal CRP. CRP has been shown in the past to have low predictive values as a single and combined biomarker for LRTI in need of antimicrobial treatment [23–25]. In addition, sensitivities and specificities for differentiation between suspected viral and bacterial LRTI have been revealed to be highly variable on several occasions [23, 24, 26–28]. Pediatric data on the impact of CRP on antibiotic prescribing for LRTI is not available to the best of our knowledge. A study in adults, however, showed a CRP exceeding 50 mg/L to be a strong driver (OR of 98.1) for antimicrobial prescription, consistent with our results [9]. Along the same lines evidence on CRP-guided reduction of antibiotic use in adult LRTI is emerging with a recent meta-analysis showing a moderate reduction of antibiotic prescription (ORs 0.30–0.73) when CRP thresholds are included in treatment decision making [29]. The second strongest association with increased antibiotic prescription was found for an elevated WBC count. This is somewhat surprising as weak predictive performance for WBC as single and combined laboratory parameter in pediatric LRTI is well known [26–28, 30–33]. This may be explained by older pediatric data showing WBC count >15 G/L to be predictive of a good response to antimicrobial treatment in pediatric pneumonia [34]. However, this study of the 1970s did not etiologically evaluate viral or mycoplasma species. Modern studies assessing a broader etiologic spectrum showed that the nasopharyngeal presence of *Mycoplasma pneumoniae*, influenza virus, or adenovirus is associated with elevated WBC as well, blurring its diagnostic performance in LRTI [31–33].

In the ProPAED study, physicians were encouraged to order WBC and CRP at study enrolment in all age groups. This might have introduced bias, as laboratory evaluation is not routinely necessary in young children. However, including only patients ≥ 2 years of age in a subanalysis did not change the result: CRP and WBC remained significantly associated with antibiotic prescribing (CRP: OR = 7.3 vs 9.3 in full analysis; WBC; OR = 6.3 vs 7.3, (all p -values < 0.01)).

Clinical parameters associated with increased antibiotic prescription

Clinical parameters predictive for antibiotic prescribing were pleuritic pain and body temperature. Pleuritic pain may be associated with more severe forms of pneumonia particularly with exudative pleural effusions or empyema, for which hospitalization and intravenous antibiotic treatment is required. However, previous studies have shown that sensitivity of pleuritic pain for severe pneumonic infection in young children is limited as children with any LRTI or non-complicated CAP can present with chest or abdominal pain without evidence of effusions or empyema [4, 35]. Further, young children may not be able to adequately locate pleuritic pain and they might rather state non-specific abdominal pain instead. Toddlers may not be able to communicate mild pleuritic pain as a symptom. Accordingly, in the sensitivity analysis, pleuritic pain lost its association with antibiotic prescribing in children younger than 2 years. In the present study population, only five of 48 patients with pleuritic pain had unilateral pleural effusions determined by expert chest radiograph review, none of them required drainage. An underestimation of the clinical sign pleuritic pain in young children cannot be excluded.

Body temperature has been shown to be a predictor of antibiotic prescribing in a large Italian study including all kinds of respiratory tract infections in pediatric primary care setting. Fever $>38^{\circ}\text{C}$ resulted in a two-fold chance of being treated with antibiotics (OR 2.34; 95%CI 1.97–2.79) [5], which is in line with our findings. Fever is included in CAP definitions and reassessment algorithms triggering thorough examination, as persistent fever may indicate development of complications and complications can predict prolonged fever [36, 37]. However, fever is common in children with LRTI on first presentation (88–96%) and non-specific to viral or bacterial etiology and thus need for antibiotic prescription [38].

Tachypnea is included in WHO, Infectious Diseases Society of America (IDSA) and British Thoracic Society (BTS) guidelines as a clinical sign for severity of LRTI disease [2, 4, 39]. It is an easily accessible clinical feature suitable for initial and follow-up evaluation of LRTI. The present analysis failed in univariate regression analysis to show tachypnea as a predictor of antibiotic prescribing. One explanation is that tachypnea as a sign of LRTI is observed in both, viral LRTI and bacterial LRTI [40]. Another explanation might be that clinicians rarely consider this parameter and more likely rely on oxygen saturation.

Incomplete pneumococcal vaccination status was associated with increased antibiotic prescription in univariate regression analyses, but not in the multivariate model. However, incomplete vaccination might drive physicians to suspect pneumococcal infection in those not vaccinated, and this may influence their prescription behavior. It may be well possible, that incomplete vaccination status might be associated with antibiotic use in future studies.

Clinical parameters associated with decreased antibiotic prescription

Wheezing and dyspnea were independently associated with lower rates of antibiotic prescribing: Particularly wheezing is associated mostly with viral infections, which is also reflected by international guidelines where antibiotic treatment is not recommended for children with mild fever, wheeze, and signs of upper respiratory tract infections [2, 4].

In multivariate analysis, the combination of six factors including CRP, dyspnea, WBC, body temperature, pleuritic pain and wheezing explained 80% of antibiotic prescribing and resulted in more than half of the patients presenting with LRTI to the emergency departments being treated with antibiotics. Importantly the present study demonstrates that there may be an opportunity for further reduction of antibiotic prescribing for febrile LRTI based on the availability of improved biomarkers and clinical parameters. Although some clinical symptoms did not show a statistically significant association with antibiotic prescription in our study (like bronchial breathing, late inspiratory crackles, elevated heart/respiratory rate, pleural rub, all with OR point estimates < 2.0), their consideration in future larger studies could still be of advantage: with our sample size we calculate a power of only $< 60\%$ to detect such a significant univariate OR ≤ 2 .

One limitation of the study is that the impact of chest radiographs on antibiotic prescribing could not be assessed due to potential selection bias. Chest radiographs were not mandatory for the initial evaluation during the ProPAED study resulting in chest radiographs being performed in only 87% of all ProPAED patients. However, a background re-assessment of the chest radiographs based on the WHO standard [41] showed a 66% diagnostic accordance with the physicians' emergency room diagnosis (S9 Table). Inter-observer agreement in the evaluation of chest radiographs in previous studies has been shown to be highly variable: depending on the study setting between 20 to 91% [42–44]. While chest radiographs were not included in the regression model, this background analysis alone demonstrates the poor accordance of chest radiographs with clinical diagnosis and consecutive treatment.

A limitation of the original ProPAED trial with a potential impact on this analysis is the treatment of the intervention group and the standard care control group by the same emergency department team. First, a Hawthorne effect for being under study observation and second, a spill-over effect for learning from the rather restrictive PCT guidance intervention group cannot not be excluded. Especially, as the antibiotic prescription rate was indeed lower in the ProPAED control group compared to historical controls from the same center [10]. The fact of being part in a prospective study with the aim to reduce antibiotic prescription might be an independent factor driving physicians to decreased antibiotic prescription.

A further limitation may be that interactions between patients, parents and physicians were not included as endpoints in the original ProPAED study. Thus, these variables could not be

included in the regression analysis. So far, large-scale studies have provided valuable insights into non-clinical factors influencing physicians when making decisions on LRTI treatment. Ahmed et al. showed that pediatric patients are at lowest risk for antibiotic exposure when visiting pediatricians in contrast to emergency department doctors or family practitioners, as the antibiotic prescribing rate in pediatricians is one-third of the rate in emergency department staff [6]. Moro et al. showed that the physicians' perception of parents expectations to receive antibiotic treatment for their child's LRTI was a strong driver for antibiotic prescribing (OR 12.8), even when this perception was wrong, as revealed in parents interviews before consultation [5]. These studies have been performed in general practice and influence of parental treatment expectations remain to be elucidated in the tertiary care setting.

Our model did explain 80% of decisions to prescribe antibiotics but leaves 20% unexplained. Several large studies as well as the latest meta-analysis of Lucas et al. demonstrated that a main factor driving antibiotic prescribing in pediatric LRTI is diagnostic uncertainty [5, 45, 46]. However, we have not assessed diagnostic uncertainty in the prescribing physician. As this may be a significant factor also in tertiary care decision making, we suggest quantification of this impact in further research.

Conclusions

CRP and WBC were strong drivers of antibiotic prescribing in children with febrile LRTI, in spite of their known poor predictive value for antibiotic requirement. In contrast, antibiotics were used prudently in patients with dyspnea and wheezing. Building on current guidelines for antibiotic treatment in children with febrile LRTI, a reliable decision algorithm for safe antibiotic withholding considering the laboratory and clinical factors evaluated in this study may have the potential to further reduce antibiotic prescribing.

Supporting information

S1 Table. Status classification for *Haemophilus influenzae* type b vaccination.
(PDF)

S2 Table. Status classification for *Pneumococcus* vaccination.
(PDF)

S3 Table. Reference ranges for white blood cell count (WBC) for university of basel children's hospital, Switzerland.
(PDF)

S4 Table. Reference ranges for white blood cell count (WBC) for Kantonsspital Aarau, Switzerland.
(PDF)

S5 Table. Classification of tachypnea for age.
(PDF)

S6 Table. Classification of tachycardia for age.
(PDF)

S7 Table. Variables with significant associations antibiotic prescribing in univariate logistic analysis.
(PDF)

S8 Table. False positive and false negative prediction of antibiotic treatment by the multivariate logistic model.

(PDF)

S9 Table. Agreement of clinical and expert chest radiography diagnosis.

(PDF)

S1 Fig. Exploratory analysis for association of respiratory rate and heart rate with antibiotic prescribing.

(PDF)

S2 Fig. Transformation of C-reactive protein for logistic regression.

(PDF)

S3 Fig. Body temperature for logistic regression.

(PDF)

S4 Fig. Days of fever for logistic regression.

(PDF)

S5 Fig. Transformation of age for logistic regression.

(PDF)

S1 Text. Example codes of the main analyses.

(PDF)

Acknowledgments

ProPAED study group membership: Arnold Amacher, Gerald Berthet, Heiner C. Bucher, Michael Buettcher, Sabine Faisst, Muriel Gambon, Juerg Hammer, Beat Mueller, Diana Reppucci, Juliane Schaefer, Jacques Schneider, Philipp Schuetz, Jody Staehelin-Massik, Daniel Trachsel, Urs B. Schaad, Edda Weimann, Urs Zumsteg. We thank Andrew Atkinson for discussions about the statistical analysis and help in the sensitivity analysis using multiple data imputation.

Author Contributions

Conceptualization: Verena Gotta, Philipp Baumann, Nicole Ritz, Aline Fuchs, Gurli Baer, Jessica M. Bonhoeffer, Ulrich Heininger, Gabor Szinnai, Jan Bonhoeffer.

Data curation: Verena Gotta, Philipp Baumann, Gurli Baer, Jessica M. Bonhoeffer, Gabor Szinnai, Jan Bonhoeffer.

Formal analysis: Verena Gotta, Philipp Baumann.

Investigation: Philipp Baumann, Gurli Baer, Jessica M. Bonhoeffer, Ulrich Heininger, Gabor Szinnai, Jan Bonhoeffer.

Methodology: Verena Gotta, Philipp Baumann, Nicole Ritz, Aline Fuchs, Gurli Baer, Jessica M. Bonhoeffer, Ulrich Heininger, Gabor Szinnai, Jan Bonhoeffer.

Project administration: Ulrich Heininger, Jan Bonhoeffer.

Resources: Philipp Baumann, Nicole Ritz, Gurli Baer, Jessica M. Bonhoeffer, Ulrich Heininger, Gabor Szinnai, Jan Bonhoeffer.

Supervision: Ulrich Heininger, Jan Bonhoeffer.

Validation: Verena Gotta, Philipp Baumann, Aline Fuchs, Gurli Baer, Jessica M. Bonhoeffer, Gabor Szinnai, Jan Bonhoeffer.

Visualization: Verena Gotta, Philipp Baumann, Aline Fuchs, Gurli Baer, Gabor Szinnai, Jan Bonhoeffer.

Writing – original draft: Verena Gotta, Philipp Baumann.

Writing – review & editing: Verena Gotta, Philipp Baumann, Nicole Ritz, Aline Fuchs, Gurli Baer, Jessica M. Bonhoeffer, Ulrich Heininger, Gabor Szinnai, Jan Bonhoeffer.

References

1. Walker CLF, Rudan I, Liu L, Nair H, Theodoratou E, Bhutta ZA, et al. Global burden of childhood pneumonia and diarrhoea. *The Lancet*. 2013; 381(9875):1405–16.
2. Bradley JS, Byington CL, Shah SS, Alverson B, Carter ER, Harrison C, et al. The Management of Community-Acquired Pneumonia in Infants and Children Older Than 3 Months of Age: Clinical Practice Guidelines by the Pediatric Infectious Diseases Society and the Infectious Diseases Society of America. *Clin Infect Dis*. 2011; 53(7):e25–e76. <https://doi.org/10.1093/cid/cir531> PMID: 21880587
3. Lemaître C, Angoulvant F, Gabor F, Makhoul J, Bonacorsi S, Naudin J, et al. Necrotizing Pneumonia in Children: Report of 41 Cases Between 2006 and 2011 in a French Tertiary Care Center. *Pediatr Infect Dis J*. 2013; 32(10):1146–9 <https://doi.org/10.1097/INF.0b013e31829be1bb> PMID: 23722529
4. Harris M, Clark J, Coote N, Fletcher P, Harnden A, McKean M, et al. British Thoracic Society guidelines for the management of community acquired pneumonia in children: update 2011. *Thorax*. 2011; 66 (Suppl 2):ii1–ii23. <https://doi.org/10.1136/thoraxjnl-2011-200598> PMID: 21903691
5. Moro ML, Marchi M, Gagliotti C, Di Mario S, Resi D. Why do paediatricians prescribe antibiotics? Results of an Italian regional project. *BMC Pediatr*. 2009; 9(1):1–9. <https://doi.org/10.1186/1471-2431-9-69> PMID: 19895678
6. Nadeem Ahmed M, Muyot MM, Begum S, Smith P, Little C, Windemuller FJ. Antibiotic Prescription Pattern for Viral Respiratory Illness in Emergency Room and Ambulatory Care Settings. *Clin Pediatr (Phila)*. 2010; 49(6):542–7. <https://doi.org/10.1177/0009922809357786> PMID: 20075029
7. Goossens H, Ferech M, Vander Stichele R, Elseviers M. Outpatient antibiotic use in Europe and association with resistance: a cross-national database study. *The Lancet*. 2005; 365(9459):579–87.
8. Koller D, Hoffmann F, Maier W, Tholen K, Windt R, Glaeske G. Variation in antibiotic prescriptions: is area deprivation an explanation? Analysis of 1.2 million children in Germany. *Infection*. 2012; 41 (1):121–7. <https://doi.org/10.1007/s15010-012-0302-1> PMID: 22826031
9. Jakobsen KA, Melbye H, Kelly MJ, Ceynowa C, Mölstad S, Hood K, et al. Influence of CRP testing and clinical findings on antibiotic prescribing in adults presenting with acute cough in primary care. *Scand J Prim Health Care*. 2010; 28(4):229–36. <https://doi.org/10.3109/02813432.2010.506995> PMID: 20704523
10. Baer G, Baumann P, Buettcher M, Heininger U, Berthet G, Schäfer J, et al. Procalcitonin Guidance to Reduce Antibiotic Treatment of Lower Respiratory Tract Infection in Children and Adolescents (Pro-PAED): A Randomized Controlled Trial. *PLoS One*. 2013; 8(8):e68419. <https://doi.org/10.1371/journal.pone.0068419> PMID: 23936304
11. Esposito S, Tagliabue C, Piccioli I, Semino M, Sabatini C, Consolo S, et al. Procalcitonin measurements for guiding antibiotic treatment in pediatric pneumonia. *Respir Med*. 2011; 105(12):1939–45. <https://doi.org/10.1016/j.rmed.2011.09.003> PMID: 21959024
12. Stocker M, Fontana M, el Helou S, Wegscheider K, Berger TM. Use of Procalcitonin-Guided Decision-Making to Shorten Antibiotic Therapy in Suspected Neonatal Early-Onset Sepsis: Prospective Randomized Intervention Trial. *Neonatology*. 2010; 97(2):165–74. <https://doi.org/10.1159/000241296> PMID: 19776651
13. Christ-Crain M, Jaccard-Stolz D, Bingisser R, Gencay MM, Huber PR, Tamm M, et al. Effect of procalcitonin-guided treatment on antibiotic use and outcome in lower respiratory tract infections: cluster-randomised, single-blinded intervention trial. *Lancet*. 2004; 363(9409):600–7. Epub 2004/02/28. [https://doi.org/10.1016/S0140-6736\(04\)15591-8](https://doi.org/10.1016/S0140-6736(04)15591-8) PMID: 14987884.
14. Christ-Crain M, Muller B. Biomarkers in respiratory tract infections: diagnostic guides to antibiotic prescription, prognostic markers and mediators. *Eur Respir J*. 2007; 30(3):556–73. <https://doi.org/10.1183/09031936.00166106> PMID: 17766633.
15. Christ-Crain M, Stolz D, Bingisser R, Muller C, Miedinger D, Huber PR, et al. Procalcitonin guidance of antibiotic therapy in community-acquired pneumonia: a randomized trial. *Am J Respir Crit Care Med*.

- 2006; 174(1):84–93. Epub 2006/04/11. <https://doi.org/10.1164/rccm.200512-1922OC> PMID: [16603606](https://pubmed.ncbi.nlm.nih.gov/16603606/).
16. Muller B, Christ-Crain M, Nylen ES, Snider R, Becker KL. Limits to the use of the procalcitonin level as a diagnostic marker. *Clin Infect Dis*. 2004; 39(12):1867–8. <https://doi.org/10.1086/426148> PMID: [15578415](https://pubmed.ncbi.nlm.nih.gov/15578415/).
17. Schuetz P, Briel M, Christ-Crain M, Stolz D, Bouadma L, Wolff M, et al. Procalcitonin to guide initiation and duration of antibiotic treatment in acute respiratory infections: an individual patient data meta-analysis. *Clin Infect Dis*. 2012; 55(5):651–62. Epub 2012/05/11. <https://doi.org/10.1093/cid/cis464> PMID: [22573847](https://pubmed.ncbi.nlm.nih.gov/22573847/)
18. Domenech VS, Nylen ES, White JC, Snider RH, Becker KL, Landmann R, et al. Calcitonin gene-related peptide expression in sepsis: postulation of microbial infection-specific response elements within the calcitonin I gene promoter. *J Investig Med*. 2001; 49(6):514–21. <https://doi.org/10.2310/6650.2001.33628> PMID: [11730087](https://pubmed.ncbi.nlm.nih.gov/11730087/).
19. Muller B, White JC, Nylen ES, Snider RH, Becker KL, Habener JF. Ubiquitous expression of the calcitonin-receptor-like receptor 1 gene in multiple tissues in response to sepsis. *J Clin Endocrinol Metab*. 2001; 86(1):396–404. Epub 2001/03/07. <https://doi.org/10.1210/jcem.86.1.7089> PMID: [11232031](https://pubmed.ncbi.nlm.nih.gov/11232031/).
20. Bernstein D. History and Physical Examination. In: Behrman R, Kliegman R, Jenson H, editors. *Nelson Textbook of Pediatrics*. 17th ed. Philadelphia: Saunders; 2004. p. 1481.
21. Fleming S, Thompson M, Stevens R, Heneghan C, Plüddemann A, Maconochie I, et al. Normal ranges of heart rate and respiratory rate in children from birth to 18 years of age: a systematic review of observational studies. *The Lancet*. 2011; 377(9770):1011–8.
22. van Buuren S, Groothuis-Oudshoorn K. mice: Multivariate Imputation by Chained Equations in R. 2011. 2011; 45(3):67. Epub 2011-12-12. <https://doi.org/10.18637/jss.v045.i03>
23. Flood RG, Badik J, Aronoff SC. The Utility of Serum C-Reactive Protein in Differentiating Bacterial from Nonbacterial Pneumonia in Children: A Meta-Analysis of 1230 Children. *Pediatr Infect Dis J*. 2008; 27(2):95–9
24. Korppi M. Non-specific host response markers in the differentiation between pneumococcal and viral pneumonia: What is the most accurate combination? *Pediatr Int*. 2004; 46(5):545–50. <https://doi.org/10.1111/j.1442-200x.2004.01947.x> PMID: [15491381](https://pubmed.ncbi.nlm.nih.gov/15491381/)
25. Don M, Francesca V, Matti K, Mario C. Differentiation of bacterial and viral community-acquired pneumonia in children. *Pediatr Int*. 2009; 51(1):91–6. <https://doi.org/10.1111/j.1442-200X.2008.02678.x> PMID: [19371285](https://pubmed.ncbi.nlm.nih.gov/19371285/)
26. Nohynek H, Valkeila E, Leinonen M. Erythrocyte sedimentation rate, white blood cell count and serum C-reactive protein in assessing etiologic diagnosis of acute lower respiratory infections in children. *Pediatr Infect Dis J*. 1995; 14(6):484–9. PMID: [7667052](https://pubmed.ncbi.nlm.nih.gov/7667052/)
27. Korppi M, Heiskanen-Kosma T, Leinonen M. White blood cells, C-reactive protein and erythrocyte sedimentation rate in pneumococcal pneumonia in children. *Eur Respir J*. 1997; 10(5):1125–9. PMID: [9163657](https://pubmed.ncbi.nlm.nih.gov/9163657/)
28. Prat C, Dominguez J, Rodrigo C, Giménez M, Azuara M, Jiménez O, et al. Procalcitonin, C-reactive protein and leukocyte count in children with lower respiratory tract infection. *Pediatr Infect Dis J*. 2003; 22(11):963–7. <https://doi.org/10.1097/01.inf.0000095197.72976.4f> PMID: [14614368](https://pubmed.ncbi.nlm.nih.gov/14614368/)
29. McDonagh M, Peterson K, Winthrop K, Cantor A, Holzhammer B, Buckley DI. Improving Antibiotic Prescribing for Uncomplicated Acute Respiratory Tract Infections. Comparative Effectiveness Review No 163 (Prepared by the Pacific Northwest Evidence-based Practice Center under Contract No 290-2012-00014-I) AHRQ Publication No 15(16)-EHC033-EF Rockville, MD: Agency for Healthcare Research and Quality. 2016. Epub 2016/02/26.
30. Korppi M, Kröger L, Laitinen M. White Blood Cell and Differential Counts in Acute Respiratory Viral and Bacterial Infections in Children. *Scand J Infect Dis*. 1993; 25(4):435–40. <https://doi.org/10.3109/00365549309008524> PMID: [8248742](https://pubmed.ncbi.nlm.nih.gov/8248742/)
31. Chen S-P, Huang Y-C, Chiu C-H, Wong K-S, Huang Y-L, Huang C-G, et al. Clinical features of radiologically confirmed pneumonia due to adenovirus in children. *J Clin Virol*. 2013; 56(1):7–12. <https://doi.org/10.1016/j.jcv.2012.08.021> PMID: [23021965](https://pubmed.ncbi.nlm.nih.gov/23021965/)
32. Peltola V, Mertsola J, Ruuskanen O. Comparison of total white blood cell count and serum C-reactive protein levels in confirmed bacterial and viral infections. *The Journal of Pediatrics*. 2006; 149(5):721–4. <https://doi.org/10.1016/j.jpeds.2006.08.051> PMID: [17095353](https://pubmed.ncbi.nlm.nih.gov/17095353/)
33. Tabain I, Ljubin-Sternak S, Cepin-Bogovic J, Markovinovic L, Knezovic I, Mlinaric-Galinovic G. Adenovirus Respiratory Infections in Hospitalized Children: Clinical Findings in Relation to Species and Serotypes. *Pediatr Infect Dis J*. 2012; 31(7):680–4. <https://doi.org/10.1097/INF.0b013e318256605e> PMID: [22517338](https://pubmed.ncbi.nlm.nih.gov/22517338/)

34. Shuttleworth DB, Charney E. Leukocyte count in childhood pneumonia. *Am J Dis Child*. 1971; 122(5):393–6. Epub 1971/11/01. PMID: [4399688](#).
35. Balfour-Lynn IM, Abrahamson E, Cohen G, Hartley J, King S, Parikh D, et al. BTS guidelines for the management of pleural infection in children. *Thorax*. 2005; 60(suppl 1):i1–i21. <https://doi.org/10.1136/thx.2004.030676> PMID: [15681514](#)
36. Lin CJ, Chen PY, Huang FL, Lee T, Chi CS, Lin CY. Radiographic, clinical, and prognostic features of complicated and uncomplicated community-acquired lobar pneumonia in children. *J Microbiol Immunol Infect*. 2006; 39(6):489–95. Epub 2006/12/14. PMID: [17164952](#).
37. Wexler ID, Knoll S, Picard E, Villa Y, Shoseyov D, Engelhard D, et al. Clinical characteristics and outcome of complicated pneumococcal pneumonia in a pediatric population. *Pediatr Pulmonol*. 2006; 41(8):726–34. <https://doi.org/10.1002/ppul.20383> PMID: [16779839](#)
38. Korppi M, Don M, Valent F, Canciani M. The value of clinical features in differentiating between viral, pneumococcal and atypical bacterial pneumonia in children. *Acta Pædiatrica*. 2008; 97(7):943–7. <https://doi.org/10.1111/j.1651-2227.2008.00789.x> PMID: [18422803](#)
39. WHO. Pocket book of hospital care for children: Second edition, Guidelines for the management of common childhood illnesses: World Health Organisation; 2013.
40. Cevey-Macherel M, Galetto-Lacour A, Gervais A, Siegrist C-A, Bille J, Bescher-Ninet B, et al. Etiology of community-acquired pneumonia in hospitalized children based on WHO clinical guidelines. *Eur J Pediatr*. 2009; 168(12):1429–36. <https://doi.org/10.1007/s00431-009-0943-y> PMID: [19238436](#)
41. Organization WH. Standardization of interpretation of chest radiographs for the diagnosis of pneumonia in children. 2001.
42. Elemraïd MA, Muller M, Spencer DA, Rushton SP, Gorton R, Thomas MF, et al. Accuracy of the Interpretation of Chest Radiographs for the Diagnosis of Paediatric Pneumonia. *PLoS One*. 2014; 9(8): e106051. <https://doi.org/10.1371/journal.pone.0106051> PMID: [25148361](#)
43. Johnson J, Kline J. Intraobserver and interobserver agreement of the interpretation of pediatric chest radiographs. *Emergency Radiology*. 2010; 17(4):285–90. <https://doi.org/10.1007/s10140-009-0854-2> PMID: [20091078](#)
44. Davies HD, Wang EE-L, Manson D, Babyn P, Shuckett B. Reliability of the chest radiograph in the diagnosis of lower respiratory infections in young children. *Pediatr Infect Dis J*. 1996; 15(7):600–4. PMID: [8823854](#)
45. Lucas PJ, Cabral C, Hay AD, Horwood J. A systematic review of parent and clinician views and perceptions that influence prescribing decisions in relation to acute childhood infections in primary care. *Scand J Prim Health Care*. 2015; 33(1):11–20. <https://doi.org/10.3109/02813432.2015.1001942> PMID: [25716427](#)
46. Horwood J, Cabral C, Hay AD, Ingram J. Primary care clinician antibiotic prescribing decisions in consultations for children with RTIs: a qualitative interview study. *Br J Gen Pract*. 2016; 66(644):e207–e13. <https://doi.org/10.3399/bjgp16X683821> PMID: [26852795](#)